



⁹⁹mTc-MDP Bone Scintigraphy in a Case of X-Linked Spondyloepiphyseal Dysplasia Tarda

X'e Bağlı Spondiloepifizyal Displazi Tarda Olgusunda ⁹⁹mTc-MDP Kemik Sintigrafisi

Ikram Zahfir, Salah Oueriagli Nabih, Meryem Aboussabr, Yassir Benameur, Omar Ait Sahel, Abderrahim Doudouh

Mohammed V Military Teaching Hospital, Clinic of Medicine and Pharmacy, Mohammed V University of Rabat Faculty of Medicine, Department of Nuclear Medicine, Rabat, Morocco

Abstract

Spondyloepiphyseal dysplasia tarda (SED) is a rare X-linked skeletal disorder affecting the spine and long bones, leading to short stature, spinal deformities, and joint stiffness. It is caused by genetic mutations, and primarily affects males. Diagnosis is confirmed by imaging and genetic testing. We report the case of a 33 years old patient with a history of X-linked SED, who presented with pain in the lower limbs and pelvis, accompanied by limited mobility. Bone scan with ⁹⁹mTc-methylene diphosphonate (MDP) revealed a polyostotic SED involvement, periarticular ossifications, and bony bridges in the active phase. The authors highlight the role of bone scan with ⁹⁹mTc-MDP in diagnosing this rare disease.

Keywords: Bone scan, ⁹⁹mTc-MDP, spondyloepiphyseal dysplasia tarda, polyostotic involvement

Öz

Spondiloepifizyal displazi tarda (SED), omurga ve uzun kemikleri etkileyen, kısa boy, omurga deformiteleri ve eklem sertliğine yol açan nadir bir X'e bağlı geçiş gösteren iskelet hastalığıdır. Genetik mutasyonlardan kaynaklanır ve öncelikli olarak erkekleri etkiler. Tanı, görüntüleme ve genetik testlerle doğrulanır. Alt ekstremitelerde ve pelviste ağrı ve hareket kısıtlılığı ile başvuran, X'e bağlı SED öyküsü olan 33 yaşında bir hastayı bildiriyoruz. ⁹⁹mTc-metilen difosfonat (MDP) ile yapılan kemik taraması, polioostotik SED tutulumunu, periartiküler ossifikasyonları ve aktif fazda kemik köprüleri gösterdi. Yazarlar, bu nadir hastalığın tanısında ⁹⁹mTc-MDP ile yapılan kemik taramasının rolünü vurgulamaktadır.

Anahtar kelimeler: Kemik taraması, ⁹⁹mTc-MDP, spondiloepifizyal displazi tarda, polioostotik tutulum

Address for Correspondence: Ikram Zahfir, Mohammed V Military Teaching Hospital, Clinic of Medicine and Pharmacy, Mohammed V University of Rabat Faculty of Medicine, Department of Nuclear Medicine, Rabat, Morocco

E-mail: ikramzahfir@gmail.com **ORCID ID:** orcid.org/0009-0002-4822-6175

Received: 17.05.2025 **Accepted:** 31.08.2025 **Epub:** 26.11.2025

Cite this article as: Zahfir I, Nabih SO, Aboussabr M, Benameur Y, Sahel OA, Doudouh A. ⁹⁹mTc-MDP bone scintigraphy in a case of X-linked spondyloepiphyseal dysplasia tarda. Mol Imaging Radionucl Ther. [Epub Ahead of Print]



Copyright© 2025 The Author. Published by Galenos Publishing House on behalf of the Turkish Society of Nuclear Medicine. This is an open access article under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND) International License.




Figure 1. A 33-year-old patient with a known history of X-linked spondyloepiphyseal dysplasia tarda (SEDT) is presenting with lower limb and pelvic pain, accompanied by limited mobility. Standard radiographs showed hump-shaped deformities in the central portion of the vertebrae, narrowed intervertebral disc spaces, and moderate epiphyseal dysplasia of the long bones. These findings were associated with bony bridges in the hips and knees.

A bone scan with ^{99m}Tc -methylene diphosphonate ^{99m}Tc -(MDP), performed 3 hours after the intravenous injection of 666 MBq (18 mCi) of ^{99m}Tc -MDP, revealed heterogeneous and abnormally increased activity with deformities of the femoral heads, elbows, knees, and ankles, a short spine, and marked thoracolumbar scoliosis, consistent with polyostotic SEDT involvement. Focal and linear areas of increased uptake were seen in the knees and hips, corresponding to periarticular ossifications and bony bridges in the active phase (Figure 1).

SED is a subgroup of bone dysplasias that affects the spine and the epiphyses of long bones. It includes three major forms: congenital SED, late-onset (tarda) SED, and SED associated with progressive arthropathy (1). More recently, at least four types of SEDT, some with autosomal recessive inheritance and others with autosomal dominant inheritance, have also been described (2). The classical form of SEDT, as illustrated in this case, is a genetic disorder inherited in an X-linked recessive pattern, manifesting exclusively in males. It results from mutations in the *SEDL* gene located on Xp22.12-p22.31 and manifests around puberty with back pain, a short stature and a short trunk, while the extremities and face remain unaffected. Early onset degenerative joint disease, particularly affecting the spine and hips, usually develops in early adulthood. Radiologic abnormalities of the vertebral bodies include platyspondyly and a central hump (1,3). Bone scan is rarely indicated in SEDT; however, the skeletal involvement observed in this condition is unusual, requiring a thorough understanding of the distribution pattern of bone abnormalities for accurate interpretation and to avoid confusion with other entities, thereby ensuring appropriate management (4).

Ethics

Informed Consent: Informed consent was obtained from the patient.

Footnotes

Authorship Contributions

Surgical and Medical Practices: I.Z., Concept: I.Z., S.O.N., M.A., A.D., Data Collection or Processing: I.Z., S.O.N., A.D., Analysis or Interpretation: I.Z., M.A., Literature Search: I.Z., S.O.N., M.A., Y.B., O.A.S., A.D., Writing: I.Z., S.O.N., A.D.

Conflict of Interest: No conflicts of interest were declared by the authors.

Financial Disclosure: The authors declare that this study has received no financial support.

References

1. Kocyigit H, Arkun R, Ozkinay F, Cogulu O, Hizli N, Memis A. Spondyloepiphyseal dysplasia tarda with progressive arthropathy. *Clin Rheumatol*. 2000;19:238-241.
2. Panda A, Gamanagatti S, Jana M, Gupta AK. Skeletal dysplasias: a radiographic approach and review of common non-lethal skeletal dysplasias. *World J Radiol*. 2014;6:808-825.
3. Gedeon AK, Tiller GE, Le Merrer M, Heuertz S, Tranebjaerg L, Chitayat D, Robertson S, Glass IA, Savarirayan R, Cole WG, Rimoin DL, Kousseff BG, Ohashi H, Zabel B, Munnich A, Gecz J, Mulley JC. The molecular basis of X-linked spondyloepiphyseal dysplasia tarda. *Am J Hum Genet*. 2001;68:1386-1397.
4. Yang F, Xu HQ, Li CL, Yang JG. Incidental finding of Tc-99m MDP bone scintigraphy in a case of X-linked spondyloepiphyseal dysplasia tarda. *Clin Nucl Med*. 2012;37:193-195.