



PET/CT Identifies Pubic Melorheostosis with High ⁶⁸Ga-PSMA Uptake in a Patient with Prostate Cancer

Prostat Kanserli Bir Hastada PET/BT ile Yüksek ⁶⁸Ga-PSMA Tutulumu Gösteren Meloreostosis Saptanması

✉ Deya Aldeen Sweedat, ✉ Ahmed Saad Abdlkadir, ✉ Akram Al-Ibraheem

King Hussein Cancer Center, Clinic of Nuclear Medicine and Positron Emission Tomography and Computed Tomography, Amman, Jordan

Abstract

Melorheostosis, a rare benign bone disorder marked by cortical thickening and irregular hyperostosis, typically affects peripheral bones but can also involve the axial skeleton. Though various molecular imaging radiotracers have been used to evaluate this condition, there are no previous reports of positive ⁶⁸Ga-prostate-specific membrane antigen positron emission tomography/computed tomography (⁶⁸Ga-PSMA PET/CT) findings in melorheostosis. This case presents a 70-year-old man with metastatic prostate cancer whose ⁶⁸Ga-PSMA PET/CT scan revealed intense ⁶⁸Ga-PSMA uptake in pubic melorheostosis seen on magnetic resonance imaging. The case highlights the possibility of ⁶⁸Ga-PSMA-avid melorheostosis and suggests ⁶⁸Ga-PSMA PET/CT may have a role in the evaluation of melorheostosis.

Keywords: Melorheostosis, ⁶⁸Ga-PSMA, diagnostic challenge, PET/CT, prostate cancer, false positive

Öz

Melorheostosis, kortikal kalınlaşma ve düzensiz hiperostosis ile karakterize nadir görülen benign bir kemik hastalığıdır. Genellikle periferik kemikleri etkiler, ancak aksiyal iskeleti de tutabilir. Bu hastalığın değerlendirilmesinde çeşitli moleküler görüntüleme radyofarmasötikleri kullanılmış olsa da, melorheostoziste pozitif ⁶⁸Ga-prostat spesifik membran antijeni pozitron emisyon tomografisi/bilgisayarlı tomografi (⁶⁸Ga-PSMA PET/BT) bulgularına dair daha önce herhangi bir bildirim bulunmamaktadır. Bu olguda, metastatik prostat kanseri olan 70 yaşındaki bir erkek hastada, manyetik rezonans görüntülemesinde saptanan pubik melorheostosis alanında ⁶⁸Ga-PSMA PET/BT incelemesinde yoğun ⁶⁸Ga-PSMA tutulumu gösterilmiştir. Bu olgu, ⁶⁸Ga-PSMA tutulumu gösteren melorheostosis olasılığını vurgulamakta ve ⁶⁸Ga-PSMA PET/BT'nin melorheostosisin değerlendirilmesinde rol oynayabileceğini düşündürmektedir.

Anahtar kelimeler: Melorheostosis, ⁶⁸Ga-PSMA, tanısal zorluk, PET/BT, prostat kanseri, yanlış pozitif

Address for Correspondence: Akram Al-Ibraheem, King Hussein Cancer Center, Clinic of Nuclear Medicine and Positron Emission Tomography and Computed Tomography, Amman, Jordan

E-mail: akramalibrahim@gmail.com ORCID ID: orcid.org/0000-0002-0978-4716

Received: 28.09.2025 Accepted: 21.04.2026 Epub: 28.04.2026 Publication Date: 06.10.2026

Cite this article as: Sweedat DA, Abdlkadir AS, Al-Ibraheem A. PET/CT identifies pubic melorheostosis with high ⁶⁸Ga-PSMA uptake in a patient with prostate cancer. Mol Imaging Radionucl Ther. 2026;35(3):260-262.



Copyright© 2026 The Author(s). Published by Galenos Publishing House on behalf of the Turkish Society of Nuclear Medicine. This is an open access article under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND) International License.




Figure 1. A 70-year-old man with metastatic hormone-sensitive prostate adenocarcinoma, treated with abiraterone and goserelin, developed progressive bilateral pubic pain. The patient provided written informed consent. Lab studies showed significantly elevated prostate-specific antigen and alkaline phosphatase levels. Pelvic X-ray revealed bony expansion with increased bone density and thickened cortex in both pubic bones (A, pelvic radiograph; arrows), which corresponded to an area of increased radiotracer uptake on skeletal scintigraphy (B, anterior and posterior planar skeletal scintigraphy; arrows). Other radiotracer-avid bone lesions were noted on skeletal scintigraphy, raising suspicion for new-onset skeletal metastases (B, anterior and posterior planar skeletal scintigraphy; arrowheads). ^{68}Ga -prostate-specific membrane antigen positron emission tomography/computed tomography (^{68}Ga -PSMA PET/CT), revealed diffuse ^{68}Ga -PSMA-avid prostatic involvement, and extensive ^{68}Ga -PSMA-avid lymphadenopathy and bone involvement [C, maximum intensity projection (MIP) PET image, D-G, axial fused images; annotations]. The pubic bones, which corresponded to areas of severe pain, displayed pronounced hyperostosis and a “dripping candlewax” sign consistent with melorheostosis; they also showed intense ^{68}Ga -PSMA uptake, with a maximum standardized uptake value of 20.2 (C, MIP PET image; H-J, axial and coronal fused PET/CT images; K-M, axial and coronal CT images; arrows). Magnetic resonance imaging further characterized these lesions (arrowheads), demonstrating bilateral cortical and subcortical hyperostosis of the pubic bones with uniformly low signal intensity on T1-weighted images (O) and predominantly low signal intensity on T2-weighted images (P), consistent with dense sclerotic bone. STIR sequences (N) revealed superimposed hyperintensity of the marrow and adjacent soft tissues, indicating associated edema and suggesting metabolically active disease. The patient was diagnosed with metastatic castration-resistant prostate cancer, was started on taxane chemotherapy, and received symptomatic therapy for melorheostosis, which led to rapid relief of pubic bone pain.

In our case, melorheostosis is located in the axial skeleton, more specifically in the pubic bones, which is a rare site of such entity (1,2). It also had increased ^{68}Ga -PSMA activity, which can be misdiagnosed as bone metastases in patients with prostate cancer (3). The observed expression pattern could be attributable to pronounced hypervascularization within melorheostotic lesions, consequently resulting in increased ^{68}Ga -PSMA expression (3,4). Also, it can be attributable to the proposition that the RAS signaling pathway, which is implicated in melorheostosis, may interact with other pathways such as TGF- β and PI3K/AKT2 (5). These pathways are known to regulate cellular processes like proliferation and differentiation, suggesting they could influence ^{68}Ga -PSMA expression in this condition (5). The differential diagnosis encompasses various osseous pathologies, including Paget’s disease, heterotopic ossification, and metastatic lesions (6,7,8). Radiological findings showing a dripping candle wax appearance indicated melorheostosis in the patient (4), highlighting the need for clinical follow-up and radiological correlation for accurate diagnosis. To the best of our knowledge, this case describes ^{68}Ga -PSMA-avid melorheostosis, reported here for the first time, and highlights its relevance as a pitfall in ^{68}Ga -PSMA PET/CT interpretation, with the aim of preventing false upstaging of prostate cancer and avoiding unwarranted treatment.

Ethics

Informed Consent: The patient provided written informed consent.

Footnotes

Authorship Contributions

Surgical and Medical Practices: D.A.S., A.S.A., A.A-I., Concept: D.A.S., A.S.A., A.A-I., Design: D.A.S., A.S.A., A.A-I., Data Collection or Processing: D.A.S., A.S.A., A.A-I., Literature Search: D.A.S., A.S.A., A.A-I., Writing: D.A.S., A.S.A., A.A-I.

Conflict of Interest: No conflicts of interest were declared by the authors.

Financial Disclosure: The authors declare that this study has received no financial support.

References

1. Kalbermatten NT, Vock P, Rüfenacht D, Anderson SE. Progressive melorheostosis in the peripheral and axial skeleton with associated vascular malformations: imaging findings over three decades. *Skeletal Radiol*. 2001;30:48-52.
2. Suresh S, Muthukumar T, Saifuddin A. Classical and unusual imaging appearances of melorheostosis. *Clin Radiol*. 2010;65:593-600.
3. de Galiza Barbosa F, Queiroz MA, Nunes RF, Costa LB, Zaniboni EC, Marin JFG, Cerri GG, Buchpiguel CA. Nonprostatic diseases on PSMA PET imaging: a spectrum of benign and malignant findings. *Cancer Imaging*. 2020;20:23.
4. Fick CN, Fratzl-Zelman N, Roschger P, Klaushofer K, Jha S, Marini JC, Bhattacharyya T. Melorheostosis: a clinical, pathologic, and radiologic case series. *Am J Surg Pathol*. 2019;43:1554-1159.
5. Yao V, Berkman CE, Choi JK, O'Keefe DS, Bacich DJ. Expression of prostate-specific membrane antigen (PSMA), increases cell folate uptake and proliferation and suggests a novel role for PSMA in the uptake of the non-polyglutamated folate, folic acid. *Prostate*. 2010;70:305-316.
6. Sachpekidis C, Bäumer P, Kopka K, Hadaschik BA, Hohenfellner M, Kopp-Schneider A, Haberkorn U, Dimitrakopoulou-Strauss A. ⁶⁸Ga-PSMA PET/CT in the evaluation of bone metastases in prostate cancer. *Eur J Nucl Med Mol Imaging*. 2018;45:904-912.
7. Jain TK, Vohra M, Sharma BS, Dhal S, Sharma LM. PSMA avidity in the heterotrophic ossification—an incidental finding on PSMA PET/CT. *Clin Nucl Med*. 2024;49:e118-e9.
8. Sasikumar A, Joy A, Nanabala R, Pillai MRA, T.A H. ⁶⁸Ga-PSMA PET/CT false-positive tracer uptake in paget disease. *Clin Nucl Med*. 2016;41:e454-e5.